

**LIST OF MEDICATION SHOULD NOT BE TAKEN PRIOR TO THE 14C-UREA BREATH TEST**

EXCLUSION PERIOD	TYPE OF MEDICATION	GENERIC#	TRADE NAME		
During Fasting and During test	Antacids		ALU-TAB DeWitts Antacid Powder Gastrogel Mylanta Titralac	Andrews Tums Antacid ENO Gaviscon Rennie	
		H2 Antagonists	Cimetidine	Magicul	Tagamet
			Ranitidine	Ausran Ranital Ulcaid Zantac	Rani 2 Ranoxyl Ultac
			Famotidine	Ausfam Pamacid Pepzan	Famohexal Pepcidine
			Nizatidine	Nizac Tazac	Tacidine
For 7 Days Prior to Test	Proton Pump Inhibitors	Lansopazole	Lanzopran Zoton	Zopral	
		Rabeprazole	Parzol Prabez Razit	Pariet Rabzole	
		Omeprazole	Acimax Meprazol Omepral Pemzo Probitor	Losec Maxor Omepro-GA Ozmep	
		Esomeprazole	Nexium		
		Pantoprazole	Gastenz Panthron Panto Pantoloc Pantomac Somac Suvacid	Ozpan Pantofast Pantozol Salpraz Sozsol Torzole	
For 14 Days Prior to Test	Cytoprotectives	Sucralfate	Carafate	Ulcyte	
For 30 Days Prior to Test	Antibiotics	Multiple actives	Nexium Hp7 Vimovo	Probitor HP7	
		Amoxicillin (Penicillins)	Abbecillin Amoxil Augmentin Cilamox Fisamox Moxidlav Ranmoxy Yomax	Alphamox Bgramin Clavulin Clamoxyl Clavycillin Curam Maxamox	
		Ampicillin (Penicillins)	Ampicyn Ibimicycn	Austrapen	
		Azithromycin (Macrolides)	Azith Zedd	Zithromax Zitrocin	
		Clarithromycin (Macrolides)	Clarac Clarithro Klacid	Clarihexal Kalixocin	
		Erythromycin (Macrolides)	EES E-Mycin	Eryc Erythrocin	
		Roxithromycin (Macrolides)	Biaxsig Roxar	Roximycin Rulide	
		Tetracyclines:	Akamin Doryx Doxsig	Doxylin Doxyhexal Minocycline#	

EXCLUSION PERIOD	TYPE OF MEDICATION	GENERIC#	TRADE NAME
			Doxy Frakas <i>Tetracycline#</i> Minomycin <i>Doxycycline#</i> Vibra-tabs
		Cephalexin	Keflex Cephatrust Ibilex Rancef
		Cefaclor (Cephalosporins)	Ceclor Karlor Keflor Ozcef
		Ciprofloxacin (Quinolones)	C-Flox Ciprol Proquin Cifran Ciproxin
	Other Anti-infectives:	Metronidazole	Flagyl Metronide Metrogyl Simplotan
		Tinidazole Rifaximin Bismuth	Fasigyn Xifaxan Denol

- # Apply to all generic brand of the active ingredient listed.
- Information compiled in this list is based on MIMS **MAY/JUNE 2013**.
- The above is **NOT** the exhausted list especially antibiotics and anti-infective drugs, if you unsure of the classification of the medication, please consult your doctor or pharmacist.
- Note: All antibiotics except Vancomycin, Sulfa (Alprim, Bactrim, Resprim, Septrin, etc) and Amphotericin B should be withheld for at least 4 weeks for maximal accuracy.